Food and Drug Administration, HHS

products adequately labeled for that use in the relevant pediatric population. Examples of how improvement might be demonstrated include, e.g., evidence of increased effectiveness in treatment, prevention, or diagnosis of disease; elimination or substantial reduction of a treatment-limiting drug reaction; documented enhancement of compliance; or evidence of safety and effectiveness in a new subpopulation; or

- (ii) The product is in a class of products or for an indication for which there is a need for additional therapeutic options.
- (d) Exemption for orphan drugs. This section does not apply to any product for an indication or indications for which orphan designation has been granted under part 316, subpart C, of this chapter.

[63 FR 66671, Dec. 2, 1998]

§ 601.28 Annual reports of postmarketing pediatric studies.

Sponsors of licensed biological products shall submit the following information each year within 60 days of the anniversary date of approval of the license to the Director, Center for Biologics Evaluation and Research:

- (a) Summary. A brief summary stating whether labeling supplements for pediatric use have been submitted and whether new studies in the pediatric population to support appropriate labeling for the pediatric population have been initiated. Where possible, an estimate of patient exposure to the drug product, with special reference to the pediatric population (neonates, infants, children, and adolescents) shall be provided, including dosage form.
- (b) Clinical data. Analysis of available safety and efficacy data in the pediatric population and changes proposed in the labeling based on this information. An assessment of data needed to ensure appropriate labeling for the pediatric population shall be included.
- (c) Status reports. A statement on the current status of any postmarketing studies in the pediatric population performed by, or on behalf of, the applicant. The statement shall include whether postmarketing clinical studies in pediatric populations were required or agreed to, and, if so, the status of

these studies shall be reported to FDA in annual progress reports of post-marketing studies under §601.70 rather than under this section.

[65 FR 59718, Oct. 6, 2000, as amended at 65 FR 64618, Oct. 30, 2000]

§ 601.29 Guidance documents.

- (a) FDA has made available guidance documents under §10.115 of this chapter to help you comply with certain requirements of this part.
- (b) The Center for Biologics Evaluation and Research (CBER) maintains a list of guidance documents that apply to the center's regulations. The lists are maintained on the Internet and are published annually in the FEDERAL REGISTER. You may request a copy of the CBER list from the Office of Communication, Training, and Manufacturers Assistance (HFM-40), Center for Biologics Evaluation and Research, Food and Drug Administration, 1401 Rockville Pike, Rockville, MD 20852-1448.

[65 FR 56480, Sept. 19, 2000]

Subpart D—Diagnostic Radiopharmaceuticals

Source: 64 FR 26668, May 17, 1999, unless otherwise noted.

§ 601.30 Scope.

This subpart applies to radiopharmaceuticals intended for in vivo administration for diagnostic and monitoring use. It does not apply to radiopharmaceuticals intended for therapeutic purposes. In situations where a particular radiopharmaceutical is proposed for both diagnostic and therapeutic uses, the radiopharmaceutical must be evaluated taking into account each intended use.

§601.31 Definition.

For purposes of this part, diagnostic radiopharmaceutical means:

(a) An article that is intended for use in the diagnosis or monitoring of a disease or a manifestation of a disease in humans and that exhibits spontaneous disintegration of unstable nuclei with the emission of nuclear particles or photons: or

§601.32

(b) Any nonradioactive reagent kit or nuclide generator that is intended to be used in the preparation of such article as defined in paragraph (a) of this section.

§ 601.32 General factors relevant to safety and effectiveness.

FDA's determination of the safety and effectiveness of a diagnostic radiopharmaceutical includes consideration of the following:

- (a) The proposed use of the diagnostic radiopharmaceutical in the practice of medicine:
- (b) The pharmacological and toxicological activity of the diagnostic radiopharmaceutical (including any carrier or ligand component of the diagnostic radiopharmaceutical); and
- (c) The estimated absorbed radiation dose of the diagnostic radiopharmaceutical.

§ 601.33 Indications.

- (a) For diagnostic radiopharmaceuticals, the categories of proposed indications for use include, but are not limited to, the following:
 - (1) Structure delineation;
- (2) Functional, physiological, or biochemical assessment:
- (3) Disease or pathology detection or assessment; and
- (4) Diagnostic or therapeutic patient management.
- (b) Where a diagnostic radiopharmaceutical is not intended to provide disease-specific information, the proposed indications for use may refer to a biochemical, physiological, anatomical, or pathological process or to more than one disease or condition.

§ 601.34 Evaluation of effectiveness.

- (a) The effectiveness of a diagnostic radiopharmaceutical is assessed by evaluating its ability to provide useful clinical information related to its proposed indications for use. The method of this evaluation varies depending upon the proposed indication(s) and may use one or more of the following criteria:
- (1) The claim of structure delineation is established by demonstrating in a defined clinical setting the ability to locate anatomical structures and to characterize their anatomy.

- (2) The claim of functional, physiological, or biochemical assessment is established by demonstrating in a defined clinical setting reliable measurement of function(s) or physiological, biochemical, or molecular process(es).
- (3) The claim of disease or pathology detection or assessment is established by demonstrating in a defined clinical setting that the diagnostic radiopharmaceutical has sufficient accuracy in identifying or characterizing the disease or pathology.
- (4) The claim of diagnostic or therapeutic patient management is established by demonstrating in a defined clinical setting that the test is useful in diagnostic or therapeutic patient management.
- (5) For a claim that does not fall within the indication categories identified in §601.33, the applicant or sponsor should consult FDA on how to establish the effectiveness of the diagnostic radiopharmaceutical for the claim.
- (b) The accuracy and usefulness of the diagnostic information is determined by comparison with a reliable assessment of actual clinical status. A reliable assessment of actual clinical status may be provided by a diagnostic standard or standards of demonstrated accuracy. In the absence of such diagnostic standard(s), the actual clinical status must be established in another manner, e.g., patient followup.

§ 601.35 Evaluation of safety.

- (a) Factors considered in the safety assessment of a diagnostic radio-pharmaceutical include, among others, the following:
 - (1) The radiation dose;
- (2) The pharmacology and toxicology of the radiopharmaceutical, including any radionuclide, carrier, or ligand;
- (3) The risks of an incorrect diagnostic determination;
- (4) The adverse reaction profile of the drug;
- (5) Results of human experience with the radiopharmaceutical for other uses; and
- (6) Results of any previous human experience with the carrier or ligand of the radiopharmaceutical when the same chemical entity as the carrier or ligand has been used in a previously studied product.